

DATA ON THE CHEMICAL COMPOSITION OF THE MUREŞ (MAROS) RIVER

ZOLTÁN HAJDU

Introduction

Knowledge of the chemical composition of the Romanian section of the Mureş River is very limited in the scientific literature (Bedő 1986, 1990, Lepši 1937a,b, Ujvári 1972). In April 1991 the Environmental Protection Agency (Tîrgu Mureş) held a scientific session that expanded our knowledge.

The chemical composition of the Mureş River is to that of the Carpathian Mountains and in the Carpathian Basin generally.

The mineral content of the water is low in the upper catchment area of the river, about 30-80 mg/l, where the right side from Giurgeului Mountain is a little bit higher. The mineral springs characterize this region with Na, Mg, Cl, SO₄, I, B ions, and natural carbondioxide content, but their effect on the Mureş is negligible. In the middle section of the river (Cîmpia Transilvaniei- Transylvanian Plateau), the mineral content changed drastically. These changes caused some effects as a (1) different type of minerals in the catchment area, where content of sodium and chloride are higher; (2) increasing of natural heavy-metal content; and (3) important human effects. The lower section is also polluted with high mineral content.

Materials and methods

The evaluation of the chemical composition of the Mureş river was made by data from the Environment Protection Agency and the University of Medicine from Tîrgu Mureş. Statistical analyses were made for evaluation.

Results and discussion

Some parameters were first compared from a section of the river near Tîrgu Mureş. Average values of the years 1975-1979 and 1986 (Table 1.), where compared, seem to show that a majority of the values increased along the river. On the basis of the comparative investigation between periods, no significant difference in parameters, such as temperature, turbidity and colour were shown. At the same time increasing of other parameters is significant at (all the) time scale. We can presume the changes were caused by human activity.

Tab.1. Comparative water analysis of the Middle-Mureş

	1975-1979	1986	Difference	%
Water temperature (gr.C)				
1.Brâncovenesci	11.1	10.4	-0.70	-7.0
2.Glodeni	12.5	11.2	-1.30	-10.4
3.Tg.Mureş	13.2	12.4	-0.60	-4.55
4.Ungheni	14.6	13.7	-0.90	-6.17
5.Cipău	15.0	15.2	+0.20	+1.34
6.Luduş	19.2	18.9	-0.30	-1.57
7.Chețani	19.3	16.5	-2.80	-14.51
Turbidity value				
1.Brâncovenesci	5.4	5.7	+0.30	+5.56
2.Glodeni	8.7	6.0	-2.70	-31.04
3.Tg.Mureş	10.6	6.9	-3.70	-34.91
4.Ungheni	15.5	9.7	-5.80	-37.42
5.Cipău	13.6	16.9	+3.30	+24.27
6.Luduş	19.2	25.4	+6.20	+32.3
7.Chețani	16.8	14.5	-2.30	-13.69
Color (Pt-Co degree)				
1.Brâncovenesci	11.8	12.3	+0.50	+4.24
2.Glodeni	13.2	10.4	-2.80	-21.22
3.Tg.Mureş	13.0	13.1	+0.10	+0.77
4.Ungheni	14.3	14.3	0	0
5.Cipău	14.8	16.6	+1.80	+12.17
6.Luduş	13.3	15.5	+2.20	+16.55
7.Chețani	14.2	14.0	-0.20	-1.41
Total dissolved solids mg/l				
1.Brâncovenesci	79.8	135.4	+45.6	+51.21
2.Glodeni	121.1	140.8	+19.7	+16.27
3.Tg.Mureş	143.7	160.4	+16.7	+11.62
4.Ungheni	213.7	220.9	+7.2	+3.37
5.Cipău	216.2	235.4	+19.2	+8.88
6.Luduş	252.3	283.1	+30.8	+12.21
7.Chețani	292.2	306.9	+14.7	+5.03
Conductivity µS/cm				
1.Brâncovenesci	125.4	180.7	+55.3	+44.1
2.Glodeni	184.4	187.7	+3.3	+1.79
3.Tg.Mureş	194.4	213.7	+21.6	+11.29
4.Ungheni	274.5	294.5	+20.0	+7.29
5.Cipău	288.0	317.2	+29.2	+10.14
6.Luduş	336.2	377.5	+41.3	+12.29
7.Chețani	389.9	389.2	-0.7	-0.18
Total suspended solids mg/l				
1.Brâncovenesci	100.8	139.4	+38.6	+38.3
2.Glodeni	125.0	149.4	+24.4	+19.52
3.Tg.Mureş	146.5	176.8	+30.3	+20.69
4.Ungheni	212.8	223.2	+10.4	+4.89
5.Cipău	218.6	244.8	+26.2	+11.99
6.Luduş	266.5	311.5	+45.0	+16.89
7.Chețani	300.9	310.7	+9.8	+3.26

Table 1. (continued)

	1975-1979	1986	Difference	%
Total hardness (G.d.)				
1.Brîncoveniști	2.9	4.7	+1.8	+62.07
2.Glodeni	3.6	7.5	+3.9	+108.34
3.Tg.Mureș	4.1	6.5	+2.4	+58.54
4.Ungheni	5.7	8.2	+2.3	+40.35
5.Cipău	6.4	9.6	+3.2	+50.0
6.Luduș	8.1	9.9	+1.8	+22.23
7.Chețani	9.3	12.0	+2.7	+29.04
Redox pot. (v)				
1.Brîncoveniști	0.374	0.447	+0.073	+19.52
2.Glodeni	0.384	0.428	+0.044	+11.46
3.Tg.Mureș	0.380	0.423	+0.043	+11.32
4.Ungheni	0.383	0.428	+0.045	+11.75
5.Cipău	0.381	0.410	+0.029	+7.62
6.Luduș	0.378	0.400	+0.022	+5.82
7.Chețani	0.379	0.401	+0.022	+5.81
pH-value				
1.Brîncoveniști	7.19	7.41	+0.22	+3.06
2.Glodeni	7.44	7.58	+0.14	+1.89
3.Tg.Mureș	7.52	7.61	+0.09	+1.20
4.Ungheni	7.17	6.64	+0.47	+6.56
5.Cipău	7.18	7.56	+0.38	+5.30
6.Luduș	7.25	7.64	+0.39	+5.38
7.Chețani	7.15	7.76	+0.61	+8.54
rH-value				
1.Brîncoveniști	27.6	30.3	+2.7	+9.79
2.Glodeni	28.1	29.7	+1.6	+5.70
3.Tg.Mureș	28.2	29.8	+1.6	+5.68
4.Ungheni	26.4	29.0	+2.6	+9.85
5.Cipău	26.6	29.1	+2.5	+9.40
6.Luduș	27.5	28.7	+1.2	+4.47
7.Chețani	27.3	28.5	+1.2	+4.40

Recent longitudinal examination of the Romanian section (Table 2.) confirm earlier information about the water quality of the Mureș. There are two important influences which fundamentally change the conditions. First, communal and industrial sewages of Tîrgu Mureș decrease the dissolved oxygen content, increase ammonium, nitrate and nitrite content. Similarly the content of macro-ions as chloride, sulphate, calcium, magnesium and sodium are enlarged (see sampling point at Ungheni). Second, the river Tîrnava transports higher chloride-, sulphate- calcium- and sodium ions, which causes changes in the water type of the Mureș, from Ca-type into Na-type (see sampling point at Mihalț).

Increases in the salt content and the load of organic materials was considerable along the river; similarly mineralization of reductive nitrogen forms (as ammonium and nitrite-ions).

Tab.2. Chemical composition of the Romanian section of the Mureş River.

Wetted output, cu.m/s	0.564	4.6	8.62	8.98	9.4	14.18	18.5	31.5	74.1	77	88	
km 4	km 70	km 133	km 1853	km 2072	km 22	km 325	km 348	km 393	km 440	km 590	km 598	km 740
Tzivonit Mulec	Slimcenn	Golden	Unghean	Cipdu	Ocna Mulec	Mihali Pod	Alba Uliua	Ghermar	Branisca	Lipova	Arad	Nedea
pH	7.3	8.5	7.56	7.5	7.3	7.2	7.5	7.4	7.41	7.8	7.95	7.8
Dissolved oxygen mg/l	8.0	8.4	7.74	7.5	7.3	7.2	7.5	7.4	7.41	7.8	7.95	7.8
BOD5 mg/l	0.96	1.64	2.84	3.51	5.45	6.29	8.2	8.2	8.39	5.1	10.02	9.73
COCD-AM mkg/l	2.72	2.51	2.24	3.53	5.18	7.9	7.8	6.77	6.77	7.01	5.22	4.16
COCD-CMg/l	1.12	1.42	2.84	4.97	50.9	87.46	638.28	357	223.86	216.63	164.6	166.3
Total suspended solids mg/l	2.10	150	164.3	280	290	490	1538	1324	607.33	586.66	485.3	460.7
Ca ⁺ ion mg/l	8.25	11.2	24.7	23.4	49.5	50.9	87.46	638.28	357	223.86	216.63	164.6
SO4 ²⁻ ion mg/l	46.6	24	21.3	32	214	235	65.95	99	72	75.84	50.7	51.3
NH4 ⁺ ion mg/l	9.5	7.3	5.7	7.3	6.9	20.8	87	24	24.3	12.96	7.46	6.64
NO2 ion mg/l	0.28	0.28	0.75	18.5	38	39.5	60	400	107	98.33	70	10.38
NO3 ion mg/l	0.034	0.034	0.125	1.17	1.17	4.987	0	0.66	1.5	13.23	10.13	1.56
Phenols mg/l	0	0	0.002	0.003	0.001	0.002	0.004	0.004	0.016	0.017	0.02	0.04
AN DNA delegerials mg/l	0	0	0.035	0.052	0.21	0.11	0.06	0.06	0.01	0.016	0.038	0.179

References

- 1.Ujvari, I. (1972): Geografia apelor României (Hydrogeography of Romania). Ed. științifică București, pp. 299-338.
2. Bedő, C. (1990): Poluarea Apelor de suprafață (Mureșul) (Contamination of the surface waters-Mureș). Scientific communication session, Tîrgu Mureș (Manuscript)
3. Bedő, C.(1986): Urmărirea gradului de autoepurare al apei rîului Mureș pe tronsonul Brîcovenesti-Chețani (Study on degree of the self-purification of the Mureș, between Brîcovenesti-Chețani). Scientific communication session,Tîrgu Mureș. (Manuscript)
4. Lepș, I. (1937a): Ghiaja, suspensiunile și spuma Mureșului-Deva (Ice, suspended material and surface foam in the Mureș at Deva). Buletinul Muzeului Județean Hunedoara.
5. Lepș, I. (1937b): Despre temperatură Mureșului-Deva (About temperature of Mureș at Deva). Buletinul Muzeului Județean Hunedoara.

Zoltán Hajdú, Focus Ecocenter, 4300-Tîrgu Mureș, O.P. 6. C.P. 620, Romania